

AB Aurigae Resolved: Evidence for Spiral Structure

Stuartt Corder, Josh Eisner, & Anneila Sargent

*Division of Physics, Mathematics & Astronomy, California Institute of Technology,
MS105-24, Pasadena, CA 91125*

sac,jae,afs@astro.caltech.edu

ABSTRACT

We obtained high angular resolution ($\sim 2''$) images of the $^{13}\text{CO}(J=1\rightarrow 0)$ line and 2.7 millimeter continuum emission, and slightly lower resolution images of $^{12}\text{CO}(J=1\rightarrow 0)$ and $\text{C}^{18}\text{O}(J=1\rightarrow 0)$ line emission toward the Herbig Ae star AB Aurigae. We resolve a circumstellar disk of diameter 780 AU (FWHM) with a velocity pattern consistent with a purely rotational disk at inclination 21.5° and position angle 58.6° . Using Keplerian disk models, we find a central source dynamical mass of $2.8 \pm 0.1 M_\odot$ and a cutoff radius of 615 AU for the ^{13}CO emission. Inclination, mass, and radius determined from ^{12}CO and C^{18}O observations agree with those values, given optical depth and abundance effects. As a result of the high angular resolution of our observations, we confirm the existence of spiral structure suggested by near-IR scattered light images and show that the spiral arms represent density contrasts in the disk.

Subject headings: circumstellar matter — stars: individual (AB Aur)

1. Introduction

Circumstellar disks often surround low ($\lesssim 2 M_\odot$) and intermediate ($\sim 2\text{--}10 M_\odot$) mass pre-main sequence objects, the T Tauri (TT) and Herbig Ae/Be (HAeBe) stars. Disk sizes are of order a few hundred AU with masses a few tenths of a solar mass or less (e.g. Beckwith & Sargent 1996; Natta, Grinin, & Mannings 2000). These values are similar to the proto-solar system (Weidenschilling 1977), suggesting that they may be the sites of planet formation. Disk temperature and density profiles, key properties for understanding how planets might emerge, have been inferred from spatially unresolved observations that rely on spectral energy distributions (SEDs) and require assumptions about disk morphology (e.g. Kenyon &

Hartmann 1987; Dullemond, Dominik, & Natta 2001). However, higher angular and spectral resolution measurements of the gas and dust in disks are critical to quantifying the profiles directly and providing a context in which to interpret unresolved observations. Spatially and kinematically resolved images enable measurement of stellar mass, disk mass, radius, inclination (i), position angle (PA), and substructure (e.g. Koerner, Sargent, & Beckwith 1993; Dutrey et al. 1998; Simon, Dutrey & Guilloteau 2001). Such observations of multiple spectral lines in DM Tau have allowed the exploration of vertical and radial disk structure (Dartois, Dutrey, & Guilloteau 2003).

The HAe star AB Aurigae at 144 pc (Perryman et al. 1997) has been studied by numerous authors at optical, IR and radio wavelengths. Most recently, Semenov et al. (2004, hereafter S04) based a detailed analysis on millimeter observations at 5-12'' resolution. While this angular resolution is insufficient to determine a stellar mass, their model fits lead to a disk radius of 400 ± 200 AU, i of $17_{-3}^{+6^\circ}$, and PA of $80 \pm 30^\circ$. This inclination agrees well with estimates of $i \lesssim 20\text{-}30^\circ$ from near-IR interferometry and scattered light (Millan-Gabet, Schloerb, & Traub 2001; Eisner et al. 2003, 2004; Grady et al. 1999; Fukagawa et al. 2004, hereafter F04). Earlier millimeter interferometry and recent mid-IR observations at angular resolution $\sim 0''.5$ suggest $i \sim 45\text{-}70^\circ$ (Marsh et al. 1995; Mannings & Sargent 1997, hereafter MS97; Liu et al. 2004).

High resolution interferometric images of circumstellar gas can determine the inclination definitively. It is likely that early studies of the AB Aur disk at moderate angular resolution were affected by spiral structure in the disk surface (F04) or large scale (~ 1000 AU) envelope emission (e.g. S04). Here we present high resolution ($2''.2$) 2.7 mm continuum and $^{13}\text{CO}(1\text{-}0)$ line images and slightly lower resolution maps of the $^{12}\text{CO}(1\text{-}0)$ and $\text{C}^{18}\text{O}(1\text{-}0)$ emission from AB Aur. In addition to constraining the disk properties and central mass, these enable us to detect and probe the spiral features seen in the images of F04.

2. Observations and Results

Observations of AB Aur were carried out between January and May of 2004 at the Owens Valley Radio Observatory (OVRO) millimeter-wave array. Six 10.4 m antennas with cryogenically cooled SIS receivers were used in five array configurations with baselines ranging from 15 to 480 m. We configured the correlator to observe the emission lines of ^{13}CO at 110.2 GHz and ^{12}CO at 115.3 GHz in 64 Hanning-smoothed channels of width 125 kHz, and the C^{18}O emission line at 109.8 GHz in 32, 250 kHz wide channels. The ^{12}CO and C^{18}O observations have maximum baselines of 103 and 120 m respectively. Simultaneously, we obtained 2.7 mm continuum measurements over 7 GHz of bandwidth with OVRO's COBRA

correlator (Hawkins et al. 2004). The quasars J0530+135 and 3C111 were observed at 20 minute intervals for phase and amplitude calibration. Absolute flux calibration, accurate to 10%, was based on measurements of Uranus and Neptune. Data reduction was carried out with MIR, an IDL-based package developed for the OVRO array by N. Scoville and J. Carpenter. Maps were made using MIRIAD (Sault, Teuben, & Wright 1995).

For AB Aur, Figure 1 shows contours of integrated intensity for ^{12}CO (left panel), ^{13}CO (center panel), and C^{18}O (right panel) emission at resolutions $4''.25$, $3''.25$, and $3''.9$, respectively overlaid on intensity-weighted mean velocity maps (color). Integrated line fluxes for ^{12}CO , ^{13}CO , and C^{18}O are 22.1 , 3.3 and $0.78 \text{ Jy km s}^{-1}$ over the velocity ranges 7.67 to 4.27 , 7.51 to 4.25 , and 7.37 to 4.63 km s^{-1} , respectively. Outside these ranges, no emission was detected above the 3σ level (360 , 105 , and 72 mJy/beam for ^{12}CO , ^{13}CO , and C^{18}O respectively). We combined our ^{13}CO dataset with that of MS97 (re-reduced using MIR) to provide greater sensitivity and found the ^{13}CO flux agrees with that of MS97, within the uncertainties. To emphasize global spatial and velocity structure, the images were restored with circular beams equal in area to the naturally weighted beams. It is immediately evident that the ^{12}CO and ^{13}CO emission is resolved and the velocity varies smoothly across the maps. Due to lower sensitivity and poorer velocity resolution, the C^{18}O image is barely resolved and the velocity pattern is sufficiently distorted to make analysis difficult.

In the left panel of Figure 2, contours of 2.7 mm continuum emission at resolution $2''.2$ are overlaid on the color image of ^{13}CO integrated intensity for AB Aur. Line and continuum emission peaks are spatially coincident to within $0''.3$. The optically thin nature of the 2.7 mm emission results in a direct relationship of disk mass to flux, $M_d = 100(F_\nu d^2 / \kappa_\nu B_\nu(T))$. Following MS97, we calculate the disk mass using this expression and the $11.5 \text{ mJy } 2.7 \text{ mm}$ flux, assuming a gas to dust mass ratio of 100 and $\kappa_\nu = 0.009 \text{ cm}^2 \text{g}^{-1}$ at 2.7 mm . Adopting a temperature of 40 K , we derive $M_d = 0.009 M_\odot$ with a factor of a few uncertainty. S04 obtained a similar value, $M_d = 0.013 M_\odot$, with a factor of 7 uncertainty. The center panel of Figure 2 displays our 2.7 mm continuum contours overlaid on the near-IR scattered light image of F04, based on which they suggest spiral structure. While the scattered light images trace small amounts of material, our 2.7 mm image demonstrates that the spiral features are massive and represent density contrasts, as the 2.7 mm flux directly traces the mass. The three most intense scattered light features are detected at least at the 3σ level of significance in 2.7 mm continuum, while the strong northern and southeastern scattered light arms are at better than the 5σ level. These northern and southeastern features are more obvious in the right panel of Figure 2. There, the symmetric central core of the continuum image, containing more than 75% of the total flux, has been subtracted to enhance the outer regions. Given the uncertainties in the symmetric source fit, the resulting northeastern and southeastern asymmetry residuals provide between 5% and 11% and 7% to 14% of the total emission,

respectively; residual peak locations are uncertain by less than $0''.4$, and emission peaks are at least at the 3σ level.

The smooth velocity gradient in the middle panel of Figure 1 suggests that material is bound to the star and executes Keplerian orbits, a hypothesis supported by S04. The top panel of Figure 3 displays the morphology of the ^{13}CO emission in velocity intervals of width 0.34 km s^{-1} . The resolution of these robustly weighted maps, $2''.2 \times 1''.8$, is almost a factor of three better than achieved by MS97 and S04, as is the signal-to-noise ratio. Continuum resolution and signal-to-noise are similarly improved. Robust weighting allows us to emphasize small scale features in individual velocity bins as opposed to global characteristics. Enhanced resolution and sensitivity enable a detailed comparison of the line emission at different velocities with that expected from models. The model consists of a flat, Keplerian disk with a single component power law emission profile and an outer radius cutoff. The emission profile was fit empirically and the emission was binned by line-of-sight velocity and convolved with the $2''.2 \times 1''.8$ beam. The observed emission was subtracted from the model and the goodness of fit was measured with a difference squared merit function. Our best fit model and 2σ clipped residuals are presented in the middle and bottom panels of Figure 3. Our interpretation relies entirely on the ^{13}CO fit. Given the poorer angular and spectral resolution of the ^{12}CO and C^{18}O observations, model fits for those lines were used as consistency checks only. The upper section of Table 1 lists the best fit parameters for each spectral line.

The best fit dynamical mass for the ^{13}CO data is $2.8 \pm 0.1 M_{\odot}$. Disk radius, inclination, and PA are 615_{-2}^{+8} AU , $21.5_{-0.3}^{+0.4^{\circ}}$ and $58.6 \pm 0.5^{\circ}$ respectively. The emission profile has form $S = 48.4(r/100)^{-1.28} \text{ mJy}$ inside a radius of 615 AU. Results for the ^{12}CO and C^{18}O are reasonably consistent. We discuss any large discrepancies below. Estimates using other methods to obtain i , PA and disk radius are shown in the lower section of Table 1. The uncertainties quoted throughout this work are purely statistical estimates, do not include possible errors in the distance, and are 3σ for our model fits and 1σ for other methods. If the velocity pattern is not Keplerian the errors on all disk parameters except radius could be substantially larger. Disk radius is more correlated with the assumed emission profile.

3. Discussion

Our resolved images of the gas and dust emission from AB Aur enable us to go beyond earlier millimeter observations at lower resolution and determine a central mass of $2.8 \pm 0.1 M_{\odot}$. This dynamical mass is only a little higher than the $2.4 \pm 0.2 M_{\odot}$ obtained from photometry and pre-main sequence track fitting (van den Ancker, de Winter, & Tjin A Djie

1998). Our dynamical model results in several local χ^2 minima but these are significantly, statistically deviant ($\geq 20\sigma$) from the global minimum value. Our best fit values of i , PA, and disk radius (see Table 1) are more accurate than those derived by S04 from lower resolution data and more complex models, and well within their uncertainties.

Large scale envelope emission, present for $r > 600$ AU (S04), may contaminate more optically thick CO emission. The best fit ^{12}CO emission profile described in Table 1 indicates structure extending to beyond 1000 AU and flux ratios of CO isotopomers show that ^{12}CO is optically thick. Given this, the high reduced χ^2 value (~ 7) for our ^{12}CO model fit is not surprising and demonstrates that a pure disk model does not fit the ^{12}CO emission. The largest deviation of ^{13}CO from the model ($\sim 4.5\sigma$) occurs in the second velocity bin (see Figure 3). The deviation is hardly significant and could be due to a number of factors including local density enhancement, a flared disk surface, etc. A detailed discussion of these is beyond the scope of this work, but there may be some evidence of disk flaring in the top panel of Figure 3 where the southern lobe of emission in bins 4 through 7 splits into two pieces. Given that the ^{13}CO observations fit the pure disk model much better than ^{12}CO and there is known contamination in the ^{12}CO emission, discrepancies between disk properties derived from ^{12}CO and ^{13}CO can be wholly attributed to this effect, increasing our confidence in the ^{13}CO values.

The model fits to mass and i based on ^{13}CO and C^{18}O emission are consistent within the errors and the difference in radii is not unexpected given the relatively low abundance of C^{18}O . Our PA determinations are consistent with most published values. Inclinations in the upper section of Table 1 are consistent with near-IR interferometry results (Millan-Gabet et al. 2001; Eisner et al. 2003, 2004) for the inner 1 AU of the disk, and with outer disk values from S04. We checked our model values by calculating i using only the separation of the extreme velocity emission and, as Table 1 shows, again find good agreement. However, i values determined from axis ratios in the visibility plane are systematically higher than the velocity based methods. Recent determinations of $i=45\text{--}65^\circ$ and $\text{PA}=30\pm 15^\circ$ by Liu et al. (2004) from mid-IR adaptive optics observations with nulling interferometry, agree less well with our model derived values, and their inclination is much closer to the larger, axis-ratio-determined values of Table 1. We believe that discrepancies between the values of inclination and position angle derived by different methods are probably due to assumptions about AB Aur symmetry.

S04 and F04 demonstrate that the assumption of symmetry is invalid, as do our millimeter observations. The spiral structure, initially suggested by F04 and demonstrated to be a density contrast through this work (see Figure 2), is a major source of asymmetry. If spiral structure is present and streaming motions along the arms are significant, the gas dis-

tribution could be tightened along the arm, resulting in foreshortening perpendicular to that direction. Whether inclination increases or decreases depends on source geometry. In either case, the PA will be biased toward the direction of the long axis of the spiral arm. Spiral structure could also affect the i and PA values determined from velocity structure, depending on the strength of the features and the optical thickness of the line; more optically thin emission features would experience greater influence from such local density enhancements. Sub-Keplerian rotation will generally result in larger inclinations for velocity determined methods with fixed mass, but a simple gas pressure aided sub-Keplerian disk is probably too simplistic an assumption if streaming motions are altering the bulk of the gas motion along the arms.

Our observations show two effects that could be due to the influence of spiral structure: 1) the model fit to the optically thin C^{18}O line leads to a significantly greater PA than does the ^{13}CO fit. The direction of change of the PA is along the strong southeastern arm as would be expected if local streaming velocities were playing a role; 2) in the left panel of Figure 2, the ^{13}CO emission spreads well beyond the boundaries of the continuum emission in the northeast and southwest but has similar extent in the southeast. If spiral structure is changing gas morphology it would cause *larger* spatially determined inclinations and PAs thereby explaining the higher inclinations estimated from axis ratios. We conclude that the range of published values of inclination and PA can be explained by the presence of substructure. The values derived from our kinematic models may be influenced by this substructure, but we do not expect the effects to be great.

The wealth of observations of AB Aur at different wavelengths and at different spatial and spectral resolutions enables analysis of this complicated circumstellar disk from the inner 1 AU (e.g. Eisner et al. 2004) to the outer envelope beyond 1000 AU (e.g. S04). Our detailed observations give an accurate dynamical mass and resolve issues about i and PA, indicating that apparent inconsistencies are due to asymmetry. Our high resolution images in 2.7 mm continuum and ^{13}CO and C^{18}O gas emission, show evidence of spiral structure. Such structure has important implications for planet formation. If it is due to instability and persists long enough or is sufficiently strong, local collapse can occur, quickly forming large planets (Boss 2002; Pickett et al. 2003 and references therein). Alternatively, if the instability dissipates or the structure is due to existing planets (Bate et al. 2003), the increase in local density may increase the cross section for collisions of planetesimals and speed up the core accretion process (Rice et al. 2004).

Research with the OVRO array is funded in part by NSF grant AST-9981546. SC is supported by an NSF Graduate Research Fellowship and JE by a Michelson Fellowship. We thank M. Fukagawa for providing her scattered light image of AB Aur, J. Carpenter for

interest and advice, and the referee, A. Natta, for constructive comments.

REFERENCES

- Bate, M. R., Lubow, S. H., Ogilvie, G. I., & Miller, K. A. 2003, *MNRAS*, 341, 213
- Beckwith, S. V. W. & Sargent, A. I. 1996, *Nature*, 383, 139
- Boss, A. P. 2002, *ApJ*, 576, 462
- Dartios, E., Dutrey, A. & Guilloteau, S. 2003, *A&A*, 399, 773
- Dullemond, C. P., Dominik, C., & Natta, A. 2001, *ApJ*, 560, 957
- Dutrey, A., Guilloteau, S., Prato, L., Duvert, G., Schuster, K., & Menard, F. 1998, *A&A*, 338, L63
- Eisner, J. A., Lane, B. F., Akeson, R. L., Hillenbrand, L. A., & Sargent, A. I. 2003, *ApJ*, 588, 360
- Eisner, J. A., Lane, B. F., Hillenbrand, L. A., Akeson, R. L., & Sargent, A. I. 2004, *ApJ*, 613, 1049
- Fukagawa, M. et al. 2004, *ApJ*, 605, L53
- Grady, C. A., Woodgate, B., Bruhweiler, F. C., Boggess, A., Plait, P., Lindler, J., Clampin, M., & Kalas, P. 1999, *ApJ*, 523, L151
- Hawkins, D. W., Woody, D. P., Wiitala, B., Fredsti, J. & Rauch, K. P. 2004, *Proc. SPIE*, 5498, 567
- Kenyon, S. J. & Hartmann, L. 1987, *ApJ*, 323, 714
- Koerner, D. W., Sargent, A. I., & Beckwith, S. V. W. 1993, *Icarus*, 106, 2K
- Liu, W. M., Hinz, P. M., Hoffmann, W. F., Brusa, G., Miller, G., & Kenworthy, M. A. 2004, *ApJ*, in press
- Mannings, V. & Sargent, A. I. 1997, *ApJ*, 490, 792
- Marsh, K. A., van Cleve, J. E., Mahoney, M. J., Hayward, T. L., & Houck, J. R. 1995, *ApJ*, 451, 777
- Millan-Gabet, R., Schloerb, F. P., & Traub, W. A. 2001, *ApJ*, 546, 358

- Natta, A., Grinin, V. P., & Mannings, V. 2000, *Protostars and Planets IV*, ed. V. Mannings, A. P. Boss, & S. S. Russell (Tucson: University of Arizona Press), 559
- Perryman, M. A. C. et al. 1997, *A&A*, 323, 49
- Pickett, B. K., Mejia, A. C., Durisen, R. H., Cassen, P. M., Berry, D. K. & Link, R. P. 2003, *ApJ*, 590, 1060
- Rice, W. K. M., Lodato, G., Pringle, J. E., Armitage, P. J., & Bonnell, I. A. 2004, *MNRAS*, 355, 543
- Sault, R. J., Teuben, P. J., & Wright, M. C. H. 1995, in *ASP Conf. Ser. 77, Astronomical Data Analysis Software and Systems IV*, ed. R. A. Shaw, H. E. Payne, & J. J. E. Hayes (San Francisco: ASP), 433
- Semenov, D., Pavlyuchenkov, Ya., Schreyer, K., Henning, Th., Dullemond, K., & Bacmann, A. 2004, *ApJ*, in press
- Simon, M., Dutrey, A. & Guilloteau, S. 2001, *ApJ*, 545, 1034
- van den Ancker, M. E., Winter, D., & Tjin A Djie, H. R. E. 1998, *A&A*, 330, 145
- Weidenschilling, S., J. 1977, *Ap&SS*, 51, 153




Fig. 1.— OVRO millimeter array images of the ^{12}CO (left panel), ^{13}CO (center panel) and C^{18}O (right panel) emission from AB Aur. Intensity-averaged velocity maps (first moment), shown in color, are overlaid with contours of the integrated intensity (zeroth moment). Contours begin at the 3σ level and are spaced by 3σ for ^{12}CO and ^{13}CO . For C^{18}O maps spacings are 2σ . The integrated ^{12}CO , ^{13}CO , and C^{18}O maps have 1σ rms of 145, 36, and 39 mJy km s $^{-1}$ respectively. Emission was clipped at least at the 4σ level in the first moment maps. Circularly restored naturally weighted equivalent beam sizes are shown in the bottom left corners.




Fig. 2.— Left panel: 2.7 mm continuum emission for AB Aur (contours) overlaid on the ^{13}CO integrated intensity image (color). The lower left portion of the map shows the continuum beam size. The ^{13}CO beam surrounds the continuum beam in the left panel. Visible ^{13}CO emission begins at 3σ level while the continuum contours begin at 3σ and increase as 2σ where 1σ is 0.36 mJy per beam. The center panel shows the same contours of this continuum emission on the scattered light image of F04. The right panel shows contours of the continuum image with a best-fit gaussian source subtracted. Contours for this panel begin at 3σ and increase as 1σ .




Fig. 3.— Red, yellow, green, blue and dark blue emission represent approximately 2, 3.5, 5.0, 6.5 and 8.5σ above the noise. The left-most images have velocity 4.27 km s^{-1} . Image velocity width is 0.34 km s^{-1} . Large tick marks define the position $\alpha = 04^h 55^m 46^s$ and $\delta = 30^\circ 33' 05''$. Right ascension ranges from 46.58^s to 45.23^s left to right and the declination extends from $32' 54.8''$ to $33' 12.3''$ bottom to top. Upper Panel: ^{13}CO observations with $2''.2$ by $1''.8$ resolution and 1σ rms noise of 35 mJy/beam . Middle Panel: Best fit Keplerian model with $\text{mass} = 2.8 M_\odot$, $i = 21.5^\circ$, and $\text{PA} = 58.6^\circ$. Bottom Panel: Residuals of the data and model clipped at 2σ .

Table 1. Best-fit Model Parameters

Line	Mass $M_{\odot}$	Radius AU	Inclination Degrees	Position Angle Degrees	Method
^{13}CO	2.8 ± 0.1	615^{+8}_{-3}	$21.5^{+0.4}_{-0.3}$	58.6 ± 0.5	Model
C^{18}O	$2.77^{+0.09}_{-0.15}$	497^{+10}_{-3}	$21.4^{+0.7}_{-0.3}$	75 ± 2	Model
^{12}CO	$3.25^{+0.13}_{-0.14}$	1060 ± 10	$32.5^{+0.4}_{-0.5}$	61 ± 3	Model
^{13}CO	2.79	N/A	24 ± 4	69 ± 8	Extreme Velocity
C^{18}O	2.79	N/A	17 ± 6	73 ± 21	Extreme Velocity
^{12}CO	2.79	N/A	26 ± 9	66 ± 25	Extreme Velocity
^{13}CO	N/A	390 ± 15	33 ± 5	85 ± 10	UV Plane
^{12}CO	N/A	500 ± 13	41 ± 3	26 ± 4	UV Plane
Dust	N/A	170 ± 17	44 ± 12	79 ± 19	UV Plane

Note. — Dust is the 2.7 mm continuum emission. Error bars are 1σ for all but the line model where errors are 3σ . Radii are HWHM values except for the line models where they represent outer edges of the disk. Emission profile for the line model was of the form $S=S_0(\frac{r}{100\text{AU}})^{-k}$ mJy for $r \leq R_{out}$. Values of (S_0, k) are $(48.4^{+1.3}_{-0.8}, 1.28 \pm 0.04)$, $(13.4^{+1.2}_{-1.9}, 1.30^{+0.06}_{-0.08})$, and $(150 \pm 2, 1.1 \pm 0.1)$ for ^{13}CO , C^{18}O , and ^{12}CO respectively.